

NONSMOOTH LYAPUNOV FUNCTIONS AND DISCONTINUOUS CARATHÉODORY SYSTEMS

Andrea Bacciotti and Francesca Ceragioli

* *Dipartimento di Matematica del Politecnico di Torino,
C.so Duca degli Abruzzi, 24 - 10129 Torino - Italy*

Abstract: Differential equations with discontinuous righthand side and solutions intended in Carathéodory sense are considered. For these systems sufficient conditions which guarantee both Lyapunov stability and asymptotic stability in terms of nonsmooth Lyapunov functions are given. An invariance principle is also proven.

Keywords: Lyapunov functions, stability, stabilizability, discontinuous control, invariance principle

1. INTRODUCTION

The interest in the study of Lyapunov-like theorems for discontinuous systems is essentially motivated by the connection between stability and stabilizability problems. Since the papers by Sussmann (1979), Artstein (1983) and Brockett (1983) it is clear that, in order to deal with general stabilization problems, discontinuous feedback laws are needed. With the introduction of discontinuous feedback laws, the theoretical problem of giving an appropriate definition of solution for a system with discontinuous righthand side, comes out. Different approaches have been followed in the literature (Filippov, 1988; Clarke *et al.*, 1997; Ancona and Bressan, 1999). Filippov solutions in particular have been used in order to deal with discontinuous stabilizability problems. For these solutions Lyapunov methods have been widely developed (see, e.g., Clarke *et al.*, 1998; Bacciotti and Rosier, 2001; Teel and Praly, 2000). Some recent papers show that the stabilization problem can be very well approached by means of Carathéodory solutions (Ancona and Bressan, 1999, 2002, 2004; Bressan, 1998; Rifford, 2002, 2003). If the notion of Carathéodory solution is accepted as a good notion of solution for discon-

tinuous systems, in particular for systems coming from discontinuous stabilization problems, it is then interesting to develop an appropriate Lyapunov theory for them. Indeed, as far as the authors know, such a generalization of Lyapunov methods, though very easy, has not been treated in the literature before. We consider nonsmooth Lyapunov functions: nonsmoothness of Lyapunov functions is in fact unavoidable when studying stability properties of discontinuous systems. In particular, Lipschitz continuous Lyapunov functions which are also nonpathological (Valadier, 1989) are considered and a notion of derivative introduced in Shevitz and Paden (1994), improved in Bacciotti and Ceragioli (1999) and then used in Bacciotti and Ceragioli (1999, 2003), is used throughout the paper. The paper consists of two main sections. In Section 2 the fundamental tools needed in order to obtain the main results are collected. In Section 3 a Lyapunov like theorem and an invariance principle are stated, proved and illustrated by means of two examples.

2. TOOLS

2.1 Carathéodory solutions

Consider the autonomous differential equation:

$$\dot{x} = f(x) \quad (1)$$

where $x \in \mathbf{R}^n$, $f : \mathbf{R}^n \rightarrow \mathbf{R}^n$.

Definition 1. Let I be an interval of \mathbf{R} . A function $\varphi : I \rightarrow \mathbf{R}^n$ is said to be a *Carathéodory solution* of (1) on I if $\varphi(t)$ is absolutely continuous and $\frac{d}{dt}\varphi(t) = f(\varphi(t))$ for almost every $t \in I$.

In the following only Carathéodory solutions are considered and they are simply addressed as solutions. Moreover the set of solutions of (1) with initial condition $x(0) = x_0$ is denoted by \mathcal{S}_{x_0} .

The vector field $f(x)$ in (1) is in general not continuous, then existence of solutions of (1) is not guaranteed by classical theorems. General sufficient conditions on the vector field $f(x)$ in order to have existence of solutions of (1) have been studied in Ancona and Bressan (1999) and Bressan (1988). Since the interest of this paper is in the study of stability properties of the system, the following basic assumption will be made throughout the paper:

(H) for any initial condition $x_0 \in \mathbf{R}^n$ at least one solution of (1) exists and all solutions are defined on the interval $[0, +\infty)$.

Definitions of stability in the case of discontinuous systems with solutions intended in Carathéodory sense do not differ from the usual ones. From now on, the origin is assumed to be an equilibrium position for the system, i.e. $f(0) = 0$.

Definition 2. System (1) is said to be

- *Lyapunov stable* at the origin if for any $\epsilon > 0$ there exists $\delta > 0$ such that for any x_0 with $\|x_0\| < \delta$ and for any $\varphi(t) \in \mathcal{S}_{x_0}$, $\|\varphi(t)\| < \epsilon$ for all $t \geq 0$;
- *locally asymptotically stable* at the origin if it is Lyapunov stable and moreover there exists $\eta > 0$ such that for any x_0 with $\|x_0\| < \eta$ and for any $\varphi(t) \in \mathcal{S}_{x_0}$, $\lim_{t \rightarrow +\infty} \varphi(t) = 0$.

2.2 Nonpathological functions and nonpathological derivative

From now on, functions considered are, besides locally Lipschitz continuous, nonpathological. We recall the definition of nonpathological function given in Valadier (1989). In the following, $\partial_C V(x)$ denotes the Clarke gradient of the function $V(x)$ at the point x (Clarke, 1983).

Definition 3. A function $V : \mathbf{R}^n \rightarrow \mathbf{R}$ is said to be *nonpathological* if for every absolute continuous function $\varphi : \mathbf{R} \rightarrow \mathbf{R}^n$ and for a.e. t , the set $\partial_C V(\varphi(t))$ is a subset of an affine subspace orthogonal to $\dot{\varphi}(t)$.

Nonpathological functions form a wide class which includes Clarke regular functions, semiconcave and semiconvex functions. Nonpathological functions can be easily handled thanks to the following proposition (Valadier, 1989).

Proposition 1. If $V : \mathbf{R}^n \rightarrow \mathbf{R}$ is locally Lipschitz continuous and nonpathological, and $\varphi : \mathbf{R} \rightarrow \mathbf{R}^n$ is absolutely continuous, then the set $\{p \cdot \dot{\varphi}(t), p \in \partial_C V(\varphi(t))\}$ is reduced to the singleton $\{\frac{d}{dt}V(\varphi(t))\}$ for almost every t .

The notion of nonpathological derivative of a map with respect to a differential equation is now introduced. This notion is analogous to the notion of set-valued derivative of a map with respect to a differential inclusion introduced in Shevitz and Paden (1994) and improved in Bacciotti and Ceragioli (1999) (note that in Bacciotti and Ceragioli (1999), Clarke regular functions instead of nonpathological functions were used; the analogous set-valued derivative for nonpathological functions has been studied in Ceragioli (2000)).

Definition 4. Let $V : \mathbf{R}^n \rightarrow \mathbf{R}$ be a Lipschitz continuous and nonpathological function and let $f : \mathbf{R}^n \rightarrow \mathbf{R}^n$. Let $A_V = \{x \in \mathbf{R}^n : \exists c \in \mathbf{R} \text{ such that } p \cdot f(x) = c \forall p \in \partial_C V(x)\}$. If $x \in A_V$, the *nonpathological derivative of the map $V(x)$ with respect to (1) at x* is the number

$$\overline{V}_f(x) = p \cdot f(x), \quad (2)$$

where p is any vector in $\partial_C V(x)$.

Note that the set A_V is dense in \mathbf{R}^n and that the set $\mathbf{R}^n \setminus A_V$ has null Lebesgue measure.

The use of the nonpathological derivative is explained by the following corollary of Proposition 1 (see also Example 1 in Section 3).

Corollary 1. Let the function $V : \mathbf{R}^n \rightarrow \mathbf{R}$ be locally Lipschitz continuous and nonpathological, and let $\varphi(t)$ be any solution of system (1). Then $\varphi(t) \in A_V$ and $\frac{d}{dt}V(\varphi(t)) = \overline{V}_f(\varphi(t))$ for almost every t .

3. RESULTS

In this section two propositions which generalize Lyapunov method to systems with discon-

tinuous righthand side and solutions intended in Carathéodory sense are proved.

Proposition 2. Let $V : \mathbf{R}^n \rightarrow \mathbf{R}$ be positive definite, locally Lipschitz continuous and non-pathological. Let A_V be defined as in Definition 4. Assume that

$$\forall x \in A_V, \quad \dot{\bar{V}}_f(x) \leq 0. \quad (3)$$

Then

- (i) system (1) is Lyapunov stable at the origin;
- (ii) if moreover there exists a function $W : \mathbf{R}^n \rightarrow \mathbf{R}$ continuous and positive definite such that $\dot{\bar{V}}_f(x) \leq -W(x)$ for all $x \in A_V$ then system (1) is locally asymptotically stable.

PROOF. (i) In order to prove Lyapunov stability of the origin it is sufficient to prove that for any solution $\varphi(t)$ of (1) the composite function $V(\varphi(t))$ is nonincreasing. According to Corollary 1, $\varphi(t) \in A_V$ for almost every t , and $\frac{d}{dt}V(\varphi(t)) = \dot{\bar{V}}_f(\varphi(t))$ is nonpositive thanks to (3). The remaining part of the proof is standard.

(ii) From (i) it follows that for all $\epsilon > 0$ there exists δ such that for any x_0 with $\|x_0\| < \delta$ and for any $\varphi(t) \in \mathcal{S}_{x_0}$ it holds $\|\varphi(t)\| < \epsilon$ for all $t \geq 0$. Next it is shown that $\lim_{t \rightarrow +\infty} \varphi(t) = 0$ for any x_0 with $\|x_0\| < \delta$ and for any $\varphi(t) \in \mathcal{S}_{x_0}$.

First it is shown that $\lim_{t \rightarrow +\infty} V(\varphi(t)) = 0$. It has already been proved in (i) that $\frac{d}{dt}V(\varphi(t))$ is nonpositive. $V(\varphi(t))$ is then nonincreasing and bounded from below and then there exists

$\lim_{t \rightarrow +\infty} V(\varphi(t)) = l$ with $l \geq 0$. Assume by contradiction that $l > 0$. Thanks to continuity of $V(x)$ and to the fact that $V(0) = 0$, there exists $0 < \mu < \epsilon$ such that if $\|x\| < \mu$ then $V(x) < l$. Let $C = \{x \in \mathbf{R}^n : \mu \leq \|x\| \leq \epsilon\}$ and let $N = \max\{-W(x), x \in C\}$. Since W is positive definite, $N < 0$. It is clear that $\mu \leq \|x_0\| < \delta \leq \epsilon$. For any $\varphi(t) \in \mathcal{S}_{x_0}$, $\varphi(t) \in C$ for all $t \geq 0$ and for almost every t , $\frac{d}{dt}V(\varphi(t)) = \dot{\bar{V}}_f(\varphi(t)) \leq -W(\varphi(t)) \leq N$. From this inequality it follows that $V(\varphi(t)) \leq V(x_0) + Nt$ and then $\lim_{t \rightarrow +\infty} V(\varphi(t)) = -\infty$, which contradicts the fact that $V(x)$ is positive definite.

Next it is shown that $\lim_{t \rightarrow +\infty} V(\varphi(t)) = 0$ implies $\lim_{t \rightarrow +\infty} \varphi(t) = 0$. The statement is proved by contradiction. Assume there exists a sequence $\{t_n\}$ with $t_n \rightarrow +\infty$ such that there exists $\sigma > 0$ such that $\|\varphi(t_n)\| \geq \sigma$ for all n . Thanks to (i), $\varphi(t)$ is bounded, then there exists a subsequence $\{t_{n_k}\}$ such that $\varphi(t_{n_k}) \rightarrow y$ for some y with $\|y\| \geq \sigma$. It follows $0 = \lim_{t_{n_k} \rightarrow +\infty} V(\varphi(t_{n_k})) = V(y) > 0$, that is a contradiction. \square

Next a LaSalle-like invariance principle is proved (see Shevitz and Paden, 1994, Ryan, 1998, Bacciotti and Ceragioli, 1999 for other versions of the invariance principle). In order to get it, some regularity for the vector field is needed.

Definition 5. A vector field $f(x)$ is said to have the *solutions closure property* if for any sequence $\{\varphi_n(t)\}$ of solutions of (1) such that $\varphi_n(t) \rightarrow \varphi(t)$ uniformly on compact subsets of \mathbf{R} , one has that also $\varphi(t)$ is a solution of (1).

Of course any continuous vector field $f(x)$ has the solutions closure property. An important class of discontinuous vector fields with the solutions closure property is the class of patchy vector fields (see Ancona and Bressan (1999)).

In the statement of Proposition 3 the notion of weakly invariant set is needed.

Definition 6. A set M is said to be *weakly invariant* for (1) if for any $x_0 \in M$ there exists $\varphi(t) \in \mathcal{S}_{x_0}$ such that $\varphi(t) \in M$ for all $t \geq 0$.

Proposition 3. Assume that the vector field $f(x)$ has the solutions closure property. Let $V : \mathbf{R}^n \rightarrow \mathbf{R}$ be positive definite, locally Lipschitz continuous and nonpathological. Let A_V be defined as in Definition 4 and assume (3). Assume that for some $l > 0$ the connected component L_l of the level set $\{x \in \mathbf{R}^n : V(x) \leq l\}$ such that $0 \in L_l$ is bounded. Let $Z_f^V = \{x \in A_V : \dot{\bar{V}}_f(x) = 0\}$ and let M be the largest weakly invariant subset of $\bar{Z}_f^V \cap L_l$. Then for any $x_0 \in L_l$ and any $\varphi(t) \in \mathcal{S}_{x_0}$

$$\lim_{t \rightarrow +\infty} \text{dist}(\varphi(t), M) = 0. \quad (4)$$

Lemma 1. If $f(x)$ has the solutions closure property, then for any x_0 and any $\varphi(t) \in \mathcal{S}_{x_0}$ the fact that the positive limit set $\Omega(\varphi)$ is nonempty implies that it is weakly invariant.

The proof of this lemma is very similar to the proof of the analogous lemma for Filippov solutions (see Filippov (1988), page 130).

PROOF. Let $x_0 \in L_l$ and let $\varphi(t) \in \mathcal{S}_{x_0}$. First of all, note that $\varphi(t)$ is bounded. In fact if $\varphi(t)$ is not bounded on one hand there exists \bar{t} such that $\varphi(\bar{t}) \notin L_l$ and on the other hand, by the proof of Proposition 2 (i), $V(\varphi(\bar{t})) \leq V(x_0) \leq l$, which is a contradiction.

Denote by $\Omega(\varphi)$ the positive limit set of $\varphi(t)$.

Since $\varphi(t)$ is bounded then $\Omega(\varphi) \neq \emptyset$ and $\lim_{t \rightarrow +\infty} \text{dist}(\varphi(t), \Omega(\varphi)) = 0$. Next it is proved that $\Omega(\varphi) \subset \bar{Z}_f^V \cap L_l$. From the fact that $\Omega(\varphi)$

is weakly invariant it will follow (4). Let $z \in \Omega(\varphi)$. Consider the composite function $V(\varphi(t))$, which is nonincreasing and bounded from below. From this fact it follows that there exists $\lim_{t \rightarrow +\infty} V(\varphi(t)) = c$ and then $V(z) = c$ for every $z \in \Omega(\varphi)$. In fact, if $z \in \Omega(\varphi)$, there exists a sequence $\{t_n\}$, $t_n \rightarrow +\infty$ such that $\lim_{n \rightarrow +\infty} \varphi(t_n) = z$ and, thanks to the continuity of $V(x)$, $V(z) = \lim_{n \rightarrow +\infty} V(\varphi(t_n)) = \lim_{t \rightarrow +\infty} V(\varphi(t)) = c$. Due to Lemma 1, $\Omega(\varphi)$ is weakly invariant, then there exists $\tilde{\varphi}(t) \in \mathcal{S}_z$ such that $\tilde{\varphi}(t) \in \Omega(\varphi)$ for all $t \geq 0$. It follows that $V(\tilde{\varphi}(t)) = c$ for all $t \geq 0$ and $\frac{d}{dt}V(\tilde{\varphi}(t)) = 0$ for all $t \geq 0$. Since $\tilde{\varphi}(t) \in A_V$ and $\tilde{V}_f(\tilde{\varphi}(t)) = \frac{d}{dt}V(\tilde{\varphi}(t)) = 0$ for almost all $t \geq 0$, $\tilde{\varphi}(t) \in Z_f^V$ for almost all $t \geq 0$. Let now $\{t_i\}$ be a sequence such that $t_i \rightarrow 0$ and $\tilde{\varphi}(t_i) \in Z_f^V$ for all i . Since $\tilde{\varphi}(t)$ is continuous it finally follows that $z = \lim_{i \rightarrow +\infty} \tilde{\varphi}(t_i) \in \overline{Z_f^V}$. \square

Proposition 2 and 3 are now illustrated by means of two simple examples.

Example 1. Consider the two-dimensional, single input, driftless system

$$\begin{cases} \dot{x} = (x^2 - y^2)u \\ \dot{y} = 2xyu \end{cases} \quad (5)$$

(the so-called Artstein's circles example). This system can not be stabilized by means of a continuous feedback. A discontinuous stabilizing feedback is

$$u(x, y) = \begin{cases} 1 & \text{if } x < 0 \\ -1 & \text{if } x \geq 0. \end{cases} \quad (6)$$

Denote by $f(x, y)$ the righthand side of the implemented system. The function $V(x, y) = \sqrt{4x^2 + 3y^2} - |x|$ satisfies the assumptions of Proposition 2. As a sum of a function of class C^1 and a concave function, $V(x, y)$ is semiconcave and then nonpathological, but it is not differentiable when $x = 0$ (the level curves are piecewise arcs of circumferences: see Figure 1). $A_V = \mathbf{R}^2 \setminus \{(x, y) : x = 0\}$ and if $(x, y) \in A_V$ then $\tilde{V}_f(x, y) = \nabla V(x, y) \cdot f(x, y) = -W(x, y)$, where $W(x, y) = \frac{4|x|^3 + 2|x|y^2 - (x^2 - y^2)\sqrt{4x^2 + 3y^2}}{\sqrt{4x^2 + 3y^2}}$ is continuous and positive definite. Assumptions of Proposition 2 are satisfied, then the implemented system is asymptotically stable. Note that in this example the set where $V(x, y)$ is not differentiable is a curve, the y -axis, which is not included in A_V , because the vector field f is transversal to it. In this case, in order to apply Proposition 2, the condition involving the nonpathological derivative needs not to be checked at all points of \mathbf{R}^2 .

Consider now the slightly modified vector fields




Fig. 1. Level curves of $V(x, y) = (4x^2 + 3y^2)^{1/2} - |x|$

$$f^+(x, y) = \begin{cases} f(x, y) & \text{if } x \neq 0 \\ (0, y) & \text{if } x = 0 \end{cases} \quad (7)$$

$$f^-(x, y) = \begin{cases} f(x, y) & \text{if } x \neq 0 \\ (0, -y) & \text{if } x = 0. \end{cases} \quad (8)$$

Note that for $x = 0$ both these vector fields are parallel to the y -axis, i.e. to the curve where $V(x, y)$ is not differentiable, and $A_V = \mathbf{R}^2$. The condition involving the nonpathological derivative needs now to be checked at all points of \mathbf{R}^2 . By computing $\tilde{V}_{f^+}(x, y)$ and $\tilde{V}_{f^-}(x, y)$ it can be easily verified that f^+ gives rise to an unstable system, while f^- gives rise to an asymptotically stable system.

Example 2. Consider the two-dimensional control system

$$\begin{pmatrix} \dot{x} \\ \dot{y} \end{pmatrix} = uA \begin{pmatrix} x \\ y \end{pmatrix} + (1 - u)B \begin{pmatrix} x \\ y \end{pmatrix}, \quad (9)$$

where

$$A = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \text{ and } B = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \quad (10)$$

and the feedback law

$$u(x, y) = \begin{cases} 1 & \text{if } (x, y) \in D \\ 0 & \text{if } (x, y) \in \mathbf{R}^2 \setminus D \end{cases} \quad (11)$$

where $D = \{(x, y) : x > 0 \text{ and } 2x < y < 3x \text{ or } x < 0 \text{ and } 3x < y < 2x\}$.

The function $V(x, y) = \frac{x^2 + y^2}{2}$ is C^∞ , nevertheless we can not use it in order to apply classical Lyapunov theorems to the implemented system due to the fact the feedback law is discontinuous. Denote by $f(x, y)$ the righthand side of the implemented system. $A_V = \mathbf{R}^2$, $\tilde{V}_f(x, y) = \nabla V(x, y) \cdot f(x, y) \leq 0$ for all $(x, y) \in \mathbf{R}^2$ and $\overline{Z_f^V} = \mathbf{R}^2 \setminus D$. The largest weakly invariant subset of $\overline{Z_f^V}$ is the origin, then the implemented system is asymptotically stable.

Differential equations with discontinuous right-hand side and solutions intended in Carathéodory sense have been considered. The interest of such equations is motivated by the fact that they naturally arise in stabilizability problems in connection with important classes of discontinuous feedback laws. For these equations sufficient conditions which guarantee both Lyapunov stability and asymptotic stability in terms of nonsmooth Lyapunov functions are proved. Moreover an appropriate version of LaSalle invariance principle is given.

REFERENCES

- Ancona, F. and A. Bressan (1999). Patchy vector fields and asymptotic stabilization. *Esaim-Cocv* **4**, 445–472.
- Ancona, F. and A. Bressan (2002). Flow stability of patchy vector fields and robust feedback stabilization. *SIAM J. Control Optim.* **5**, 1455–1476.
- Ancona, F. and A. Bressan (2004). Stability rates for patchy vector fields. *Esaim-Cocv* **10**, 168–200.
- Artstein, Z. (1983). Stabilization with relaxed controls. *Nonlinear Analysis* **7**, 1163–1173.
- Bacciotti, A. and F. Ceragioli (1999). Stability and stabilization of discontinuous systems and nonsmooth Lyapunov functions. *Esaim-Cocv* **4**, 361–376.
- Bacciotti, A. and F. Ceragioli (2003). Nonsmooth optimal regulation and discontinuous stabilization. *Abstract and Applied Analysis* **20**, 1159–1195.
- Bacciotti, A. and L. Rosier (2001). *Liapunov Functions and Stability in Control Theory*. Springer. London.
- Bressan, A. (1998). Singularities of stabilizing feedbacks. *Rendiconti del Seminario Matematico dell’Università e del Politecnico di Torino* **4**, 87–104.
- Brockett, R. (1983). Asymptotic stability and feedback stabilization. In: *Differential Geometric Control Theory* (R. Millman R. Brockett and H. Sussmann, Eds.). pp. 181–191. Birkhäuser. Boston.
- Ceragioli, F. (2000). Discontinuous ordinary differential equations and stabilization. Ph.D. Thesis, Università di Firenze.
- Clarke, F.H. (1983). *Optimization and Nonsmooth Analysis*. Wiley and Sons. New York.
- Clarke, F.H., Yu. S. Ledyaev E.D. Sontag and A.I. Subbotin (1997). Asymptotic controllability implies feedback stabilization. *IEEE Trans. Automat. Control* **42**, 1394–1407.
- Clarke, F.H., Yu. S. Ledyaev R.J. Stern and P.R. Wolenski (1998). *Nonsmooth Analysis and Control Theory*. Springer. New York.
- Filippov, A.F. (1988). *Differential Equations with Discontinuous Right Hand sides*. Kluwer. Dordrecht.
- Rifford, L. (2001). On the existence of nonsmooth control Lyapunov functions in the sense of generalized gradients. *Esaim-Cocv* **6**, 593–611.
- Rifford, L. (2002). Semiconcave control-Lyapunov functions and stabilizing feedbacks. *SIAM J. Control Optim.* **41**, 659–681.
- Rifford, L. (2003). Singularities of viscosity solutions and the stabilization problem in the plane. *Indiana Univ. Math. J.* **52**, 1373–1396.
- Ryan, E.P. (1998). An integral invariance principle for differential inclusions with applications in adaptive control. *SIAM J. Control Optim.* **36**, 960–980.
- Shevitz, D. and B. Paden (1994). Lyapunov stability theory of nonsmooth systems. *IEEE Transaction on Automatic Control* **39**, 1910–1914.
- Sontag, E.D. (1999). Stability and stabilization: discontinuities and the effect of disturbances. In: *Nonlinear Analysis, Differential Equations and Control* (Clarke F.H. and R.J. Stern, Eds.). pp. 551–598. Kluwer. Dordrecht.
- Sussmann, H. (1979). Subanalytic sets and feedback control. *J. Differential Equations* **31**, 371–393.
- Teel, A.R. and L. Praly (2000). A smooth Lyapunov function from a class- \mathcal{KL} estimate involving two positive semidefinite functions. *Esaim-Cocv* **5**, 313–368.